

Original Paper

Development of a Mobile Health Snacktivity App to Promote Physical Activity in Inactive Adults (SnackApp): Intervention Mapping and User Testing Study

James P Sanders^{1,2}, PhD; Kajal Gokal^{1,2}; Jonah J C Thomas^{1,2}; Jonathan C Rawstorn³; Lauren B Sherar^{1,2,4}, PhD; Ralph Maddison³; Colin J Greaves⁵, PhD; Dale Eslinger^{1,2,4}; Amanda J Daley^{1,2}, PhD; Snacktivity Investigators⁶

¹Centre for Lifestyle Medicine and Behaviour, Loughborough University, Loughborough, United Kingdom

²School of Sport, Exercise and Health Sciences, Loughborough University, Loughborough, United Kingdom

³Institute of Physical Activity and Nutrition, School of Exercise and Nutrition Science, Deakin University, Burwood, Austria

⁴National Institute for Health Research Leicester Biomedical Research Centre, Leicester, United Kingdom

⁵School of Sport, Exercise and Rehabilitation Sciences, University of Birmingham, Birmingham, United Kingdom

⁶See Acknowledgments

Corresponding Author:

James P Sanders, PhD

Centre for Lifestyle Medicine and Behaviour

Loughborough University

Sir John Beckwith Building

Epinal way

Loughborough, LE11 3TU

United Kingdom

Phone: 44 01509 ext 225454

Email: J.Sanders2@lboro.ac.uk

Abstract

Background: Despite the unequivocal evidence demonstrating the benefits of being physically active, many people do not meet the recommended guidelines of at least 150 minutes of moderate- to vigorous-intensity physical activity per week. This can be changed with the development and implementation of innovative interventions. The use of mobile health (mHealth) technologies has been suggested as a mechanism to offer people innovative health behavior change interventions.

Objective: This study aims to outline the systematic, theory-driven processes and user testing applied to the development of a smartphone-based physical activity app (SnackApp) to promote participation in a novel physical activity intervention called Snacktivity. The acceptability of the app was explored and reported.

Methods: Intervention mapping involves a 6-step process, the first 4 of which were presented in this study. These steps were used to develop the SnackApp for use within the Snacktivity intervention. The first step involved a needs assessment, which included composing an expert planning group, patient and public involvement group, and gathering the views of the public on Snacktivity and the public perception of the use of wearable technology to support Snacktivity. This first step aimed to determine the overall purpose of the Snacktivity intervention. Steps 2 to 4 involved determining the intervention objectives, the behavior change theory and techniques on which the intervention is based, and the development of the intervention resources (ie, SnackApp). After the completion of steps 1 to 3 of the intervention mapping process, the SnackApp was developed and linked to a commercial physical activity tracker (Fitbit Versa Lite) for the automated capture of physical activity. SnackApp includes provisions for goal setting, activity planning, and social support. Stage 4 involved users (inactive adults, N=15) testing the SnackApp for 28 days. App engagement (mobile app use analytics) was analyzed to determine app use and to inform the further development of SnackApp.

Results: Over the study period (step 4), participants engaged with SnackApp an average of 77 (SD 80) times. On average, participants used the SnackApp for 12.6 (SD 47) minutes per week, with most of the time spent on the SnackApp dashboard and engaging, on average, 14 (SD 12.1) times, lasting 7 to 8 minutes per week. Overall, male participants used the SnackApp more than female participants did. The app rating score was 3.5 (SD 0.6) out of 5, suggesting that SnackApp was rated as fair to good.

Conclusions: This study outlines and reports data regarding the development of an innovative mHealth app using a systematic, theory-driven framework. This approach can guide the development of future mHealth programs. User testing of the SnackApp

suggested that physically inactive adults will engage with the SnackApp, indicating its applicability of use in the Snacktivity physical activity intervention.

(*JMIR Form Res* 2023;7:e41114) doi: [10.2196/41114](https://doi.org/10.2196/41114)

KEYWORDS

intervention development; physical activity; mobile health; mHealth; health app; user testing; intervention; behavior; smartphone; app; social; user; engagement; development; testing; mobile phone

Introduction

Background

The global population has become less physically active and more sedentary, and both behaviors are independently associated with poorer health outcomes and mortality [1-3]. The last two decades have seen a rapid increase in well-designed and robustly evaluated physical activity trials [4,5]. This has helped national governments and the World Health Organization to generate evidence-based physical activity recommendations [6,7]. In many countries, public health guidance recommends that adults should perform at least 150 to 300 minutes of moderate-intensity physical activity in bouts of any duration, as well as perform muscle-strengthening activities for at least 2 days a week [7]. However, few people achieve this amount of physical activity each week, and evidence suggests that initial increases in physical activity levels to meet guidelines are often not maintained over the long term [8-10]. Of particular concern are data from the United States and the United Kingdom that suggest that only 1% to 16% of adults participate in muscle-strengthening-based physical activities each week [11,12], which are important for reducing the risk of falls, fractures, and osteoporosis [13]. Another concern is the amount of time people spend being sedentary, with adults spending approximately 60% to 70% of their waking hours sitting [14]. For inactive adults, high levels of sedentary time have been associated with an increased risk of developing noncommunicable diseases (eg, type 2 diabetes) and all-cause mortality [15]. Together, this suggests that there is a need for more innovative interventions to encourage the population to regularly engage in physical activity and spend less time sedentary each day.

An alternative integrative approach to promoting physical activity that could engage and motivate people to be more physically active and break up sedentary time is a concept called *Snacktivity* [16]. Rather than emphasizing at least 150 minutes per week of physical activity, *Snacktivity* focuses on promoting short but frequent bouts of moderate- to vigorous-intensity physical activity (MVPA; known as *activity snacks*) throughout the day, such that people accumulate at least 150 MVPA minutes over the week. An *activity snack* lasts between 2 and 5 minutes, for example, walking during coffee breaks, squats while waiting for the kettle to boil, lunges while vacuuming, or getting off the bus a stop early or parking further away [16]. Other types of short-bout physical activity interventions already exist such as high-intensity interval training [17]; however, *Snacktivity* differs in purpose, duration, and intensity as an intervention approach. High-intensity interval training tends to be approximately 60 to 90 seconds in duration and at a physical intensity close to

100% exertion. In contrast, the *Snacktivity* approach aims to encourage the public to perform activity snacks in 2- to 5-minute bouts at moderate to vigorous intensity. *Snacktivity* is a multicomponent intervention approach that also encourages the public to incorporate or embed *Snacktivity* into everyday activities. *Snacktivity* is also designed to be an accessible approach to promoting physical activity in all populations, including individuals who are disabled and those who are living with chronic disease and or conditions, pregnant individuals, and older people.

Mobile Health Interventions to Promote Physical Activity in the Population

To achieve a population-level increase in physical activity levels, we need effective interventions with high reach. Given their ubiquitous use, mobile health (mHealth) interventions can reach many people, and they are relatively inexpensive to deliver and can be offered at a convenient time and place [18]. Therefore, these tools can be used to promote physical activity interventions such as *Snacktivity* in the population [19]. Furthermore, mHealth interventions can integrate key health behavior change strategies, including self-monitoring, goal setting, prompts or nudges, and feedback on behavior, which are effective in increasing participation in physical activity [20]. Evidence from systematic reviews supports the use of mHealth interventions to promote health behavior changes [18,21,22], although trials with long-term follow-up (eg, >6 months) are still required [23,24].

Intervention Mapping

Despite mHealth technologies becoming increasingly accessible, the development strategies and processes used to create interventions using them remain unclear. Appropriate steps are essential when developing evidence- and theory-based mHealth interventions to assist with effective implementation in real-world settings [25]. mHealth interventions must use a common language and a systematic and robust development process [20], including the use of theoretical frameworks (such as the intervention mapping [IM] framework) [26] to ensure that interventions are theoretically sound and can be reproduced and scaled up.

Aims

This study aimed to outline the systematic, theory-driven processes (using IM) undertaken to develop a smartphone-based physical activity app (called *SnackApp*), which is linked to a wearable consumer physical activity tracker (Fitbit) to promote participation in *Snacktivity*. In addition, the acceptability of the app as measured by the level of user engagement with the *SnackApp* was explored and reported.

Methods

Overview

Following the UK Medical Research Council's guidance for developing complex health care interventions [25,27], we used a systematic, evidence-informed approach to develop SnackApp. The development of SnackApp is part of a larger research program to test the effectiveness of Snackivity in increasing physical activity across the day in adults [16]. In brief, the SnackApp (and accompanying consumer physical activity tracker) aims to support users to incorporate Snackivity into their day through self-monitoring, by providing feedback on the number of activity snacks they have completed, by offering

users a method to set goals for Snackivity, and by creating action plans for completing activity snacks each day.

SnackApp Development

IM Framework [26] is a 6-step framework, often applied to guide the development of health behavior change interventions. The first four stages of the IM development process (which are presented in this paper) include (1) conducting a needs assessment, (2) specifying intervention outcomes and objectives, (3) designing the intervention and applying theory, and (4) refining intervention development. The final 2 steps for SnackApp development (adoption, implementation, and evaluation plan) are currently being researched [28] (Figure 1).

Figure 1. SnackApp development—intervention mapping overview. Steps 5 and 6 are ongoing [28]. MVPA: moderate- to vigorous- intensity physical activity; PPI: patient and public involvement.

Step 1	Needs assessment <ul style="list-style-type: none"> • Scoping review of the literature • Establish planning group • Stakeholder involvement and PPI • Views of the public
Step 2	Program objective <ul style="list-style-type: none"> • State the outcomes • Specify the performance objectives • Specify the determinants of behavior (eg MVPA) • Define the behavior change techniques • Define how these change techniques will be implemented into the app
Step 3	Theory-based intervention methods and practical applications <ul style="list-style-type: none"> • Define the underlying theory and processes of change
Step 4	Intervention Development—SnackApp <ul style="list-style-type: none"> • Operationalize strategies into plans – that is design and develop SnackApp • User testing
Step 5	Adoption and implementation of plan
Step 6	Evaluation Plan

Step 1: Needs Assessment

Overview

As part of the initial steps to develop SnackApp, the research team reviewed relevant literature on behavior change theory and techniques related to physical activity promotion and mHealth interventions. This helped us guide the content (ie, key behavioral strategies and techniques, app implementation, and engagement) and focus of SnackApp. The development of the SnackApp coincided with the 2019-2020 update of UK physical activity guidelines, which removed the caveat suggesting that bouts of MVPA lasting ≥ 10 minutes were needed

to derive health benefits (ie, acknowledging that any duration of MVPA is indeed health enhancing) [6].

Following this change in the physical activity guidelines and based on experimental evidence indicating that short bouts of MVPA may improve indicators of health [13,29-31], a consensus was reached from the planning group (see *Establish and Work with a Planning Group* section) that the SnackApp should aim to promote short bouts of MVPA (2-5 min), in line with the behavioral principles of Snackivity, such that users accumulate their weekly total minutes of MVPA through the accumulation of activity snacks.

Establish and Work With a Planning Group

A multidisciplinary team involving 16 individual psychologists, experts in physical activity and digital health, behavioral scientists, medical sociologists, and health care professionals was established to provide their expertise in the development and content of the SnackApp and to agree on the physical activity behaviors to be measured and the information to be feedback to users. A total of 33 meetings between some or all members of the planning group were held between September 2019 and December 2020. During these meetings, the expert planning group undertook a series of tasks described below in step 3 (eg, MoSCoW [Must Have, Should Have, Could Have, and Won't Have] framework) to determine the content, design, and behavior change techniques (BCTs) within the SnackApp.

Stakeholder and Patient and Public Involvement

During the early development of the Snacktivity intervention, 165 members of the public responded to a questionnaire regarding the concept of Snacktivity and the use of digital technology as a mechanism for behavior change. Furthermore, 3 focus groups (n=10; 6/10, 60% female) were conducted to explore the concept of Snacktivity. The focus group attendees discussed the importance of embracing digital technologies to enable people to lead healthier lifestyles. One theme that was generated related to the concept that it is necessary to ensure that there is sufficient opportunity for members of the public to be a part of the development of the SnackApp and test it before it is used. Focus group participants described the app that they were using to increase their physical activity, and they wanted to ensure that the SnackApp incorporated the best features. These included motivational messages to encourage people to engage in more activity snacks and graphics showing their achievements (unpublished data). Furthermore, throughout the planning process, we worked closely with a group of public contributors (public advisory group [PAG]; N=11) who provided feedback on the development of SnackApp and the wider Snacktivity intervention [32].

View From the Public

Following the initial stakeholder and PAG feedback stage, a survey to further assess the views of the public about Snacktivity was completed by 754 members of the public in the West Midlands [33]. When the participants were asked what would help them to perform Snacktivity throughout the day, the most frequently selected options included seeing how much Snacktivity had been completed each day, having enough examples of activity snacks to do, and receiving an alert reminding them to do Snacktivity. Furthermore, when asked about physical activity trackers and mHealth app use, most participants (678/754, 89.9%) owned a smartphone, and of them,

45% (305/678) used their mobile phones to track their physical activity. Of this 45% population, 52.1% (159/305) downloaded an app to the phone to aid in tracking their physical activity, and the most appealing features were considered to be the option to track steps and the ability to view a summary of their data. The most commonly owned physical activity tracker was the Fitbit. The 3 most appealing features of an activity tracker were notifications to move when sedentary for too long, the ability to track physical activity, and the ease of use. On the basis of these findings, the expert group agreed that the intervention should aim to increase MVPA and include both an mHealth app and the Fitbit physical activity tracker for the Snacktivity intervention.

Step 2: Specification of Outcomes, Performance Objectives, and Change Techniques

The second IM step involved developing the specific health BCTs and aligning the Snacktivity behaviors to BCTs in inactive adults. Informed by the findings from stage 1 (needs assessment), the planning group identified 3 key performance objectives of SnackApp. This included the objectives to be achieved by individuals so that they complete Snacktivity and MVPA (eg, engagement with the SnackApp) and the behavioral determinants and techniques that would influence their physical activity behavior (Table 1).

The overall purpose of SnackApp is to facilitate an increase in users' total minutes of participation in MVPA each week by encouraging engagement with Snacktivity. The three performance objectives (the behavioral outcomes intended to be achieved that would promote participation in Snacktivity and the achievement of goals) for the SnackApp or consumer physical activity tracker were as follows:

1. Education on the principles of Snacktivity and why and how it could be important for health
2. Facilitation and sustainable engagement with the SnackApp
3. Promotion of participation in minutes of MVPA per day per week

Drawing from the literature and the behavior change taxonomy [34], we aimed to achieve these 3 performance objectives through the implementation of several BCTs within the SnackApp and Fitbit physical activity tracker [34]. The BCTs were incorporated into the SnackApp and aimed to increase modifiable determinants of MVPA (eg, self-efficacy) to achieve performance objectives. The BCTs included self-monitoring, goal setting, prompts or cues, and feedback (Table 1). With the performance objectives and outcomes decided, it was important to determine how these outcomes could be changed. The next step was to determine the BCTs that the literature had suggested as effective in improving physical activity behavior.

Table 1. Program outcomes, performance objectives, and behavior change techniques (BCTs) used in the development of the SnackApp.

Performance objective, modifiable determinants, and BCT	Snacktivity intervention outcome—increase in weekly minutes of MVPA ^a (how BCTs were implemented into the app)
Learn about Snacktivity and why and how it is important for health	
Knowledge	
Provide information on the behavior	<ul style="list-style-type: none"> Written information on the health benefits of Snacktivity is provided in the form of articles on the SnackApp
Provide information on the consequences	<ul style="list-style-type: none"> Text on benefits of physical activity for health as a resource within the SnackApp
Salience of consequences	<ul style="list-style-type: none"> Emphasize the detrimental health consequences of physical inactivity via push notifications and written content in the app
Engagement with the SnackApp	
Intention	
SnackApp personalization ^b	<ul style="list-style-type: none"> Users engage with SnackApp functions tailored to their needs. <ul style="list-style-type: none"> Automatic and tailored goal setting based on their previous MVPA behavior Personalized the metrics in which the users are interested. A choice between the number of activity snacks, active minutes, and steps
Credible source	<ul style="list-style-type: none"> Users will know the SnackApp has been created by health behavior change experts
Increase daily MVPA	
Self-efficacy	
Prompt specific goal setting ^c	<ul style="list-style-type: none"> The user is prompted to use the goal-setting function in the SnackApp to formulate their own goals related to steps, active minutes, and activity snacks
Graded task ^c	<ul style="list-style-type: none"> Users can use automatic goal-setting functionality within the app, which will automatically increase or decrease user goals based on previous data as a substitute for manual goal setting
Prompt barrier identification ^d	<ul style="list-style-type: none"> Users can create action plans within the app about how to complete activity snacks throughout the day
Prompt review of behavioral goals ^c	<ul style="list-style-type: none"> Regular review of progress and increase goal for duration and bouts of physical activity by the user
Behavior instruction or demonstration	<ul style="list-style-type: none"> Exposure to modeled behavior. The SnackApp has examples of activity snacks, with videos and instructions on how they should be performed
Individualized feedback on MVPA and steps and activity snacks ^c	<ul style="list-style-type: none"> Numerical feedback (graph and push numbers) on the SnackApp will provide data on users' Snacktivity behavior
Focus on past success ^c	<ul style="list-style-type: none"> Historical graphical data displaying instances when users successfully increased their physical activity
Self-monitoring of behavior ^c	<ul style="list-style-type: none"> Encourage monitoring of own behavior, encourage users to check daily whether they have reached their Snacktivity goals via push notifications
Prompts and cues ^{c,d}	<ul style="list-style-type: none"> Prompts or push notifications for the individual to break up prolonged periods of inactivity or sedentary behavior as measured by the activity tracker
Habit formation ^d	<ul style="list-style-type: none"> Provision for users to anchor planned activity snacks to existing behaviors (eg, when I brush my teeth, I will do squats for 2 minutes)
Social reward ^c	<ul style="list-style-type: none"> Congratulatory push notification on reaching a goal
Reframing ^d	<ul style="list-style-type: none"> Framing motivational messages such as “some is good more is better”
Relapse prevention ^d	<ul style="list-style-type: none"> Provide motivational messages to promote MVPA after a period of relapse

Performance objective, modifiable determinants, and BCT	Snacktivity intervention outcome—increase in weekly minutes of MVPA ^a (how BCTs were implemented into the app)
Subjective norms	
Social support ^{c,d}	<ul style="list-style-type: none"> Provision of a forum within the app for individuals to discuss their Snacktivity
Social comparison	<ul style="list-style-type: none"> Optional provision for users to see how their peers are performing in comparison with themselves
Enjoyment	
Offer users an option of performing Activity Snacks they enjoy ^b	<ul style="list-style-type: none"> Users to choose from a library of activity snacks that they enjoy

^aMVPA: moderate- to vigorous-intensity physical activity.

^bThese are the BCTs, which were determined by the expert planning group based on their expertise and the Patient Advisory Group from their experiences.

^cThese BCTs are related to control theory and self-regulation theory.

^dThese BCTs are derived from habit formation model.

Step 3: Theory-Based Intervention Methods and Practical Applications

Overview

Step 3 of the IM process involved the selection of BCTs targeting each modifiable determinant identified in Table 1. For each determinant (ie, knowledge, self-efficacy, and enjoyment) of each performance objective (eg, increasing daily MVPA per minute), BCTs that were likely to alter the determinant (eg, self-monitoring) were selected [26]. This stage was based on the existing BCT taxonomy [34]. To ensure that different perspectives were included in this development stage, the PAG was asked, through a series of workshops, to comment on the strategies that they had found to be successful in encouraging them to be more physically active and were asked to review the BCTs that were planned to be included in the SnackApp. This increased the likelihood that they would be feasible and acceptable for potential users. In addition, as part of the process of determining which BCTs should be included, we used a MoSCoW method exercise with the PAG and the expert planning group. Briefly, MoSCoW stands for “Must have,” “Should have,” “Could have,” and “Won’t have,” and is a prioritization technique used in software development to reach a common understanding with stakeholders on the importance they place on the BCT components and strategies to be included. Members of the expert planning group were provided with a series of BCTs and how they could be implemented into the SnackApp and were asked to complete the SnackApp MoSCoW template. Responses that were deemed “Must” or “Should have” BCTs were incorporated into the SnackApp.

Underlying Theory or Process of Change

The SnackApp is based on several theoretical perspectives. The key principles in the development of SnackApp were to encourage user learning from their experience through engagement in self-regulation activities (eg, self-monitoring, feedback, and goal setting), in line with control theory [35] and self-regulation theory [36]; build intrinsic motivation; and promote autonomy (also consistent with self-determination theory) [37]. An important aim of SnackApp is to promote the formation of daily habits for Snacktivity in line with the habit

formation model [38] to maximize the potential for the maintenance of Snacktivity and physical activity. With the BCTs decided, the next step focused on the inclusion and implementation of the BCTs within Snacktivity intervention resources, in this case, the SnackApp.

Step 4: Intervention Development—SnackApp

Overview

Step 4 of the IM process was to develop the SnackApp. This step involved translating the BCTs (and the underlying processes of change) into specific SnackApp functionalities (eg, how to measure Snacktivity, how to provide feedback and goal setting, and what information about Snacktivity should be included in the SnackApp). The planning group discussed and guided the scope, selection of behavioral strategies (ie, how BCTs would be translated into app features), and the feasibility of encouraging Snacktivity via an app. To facilitate the initiation and adoption of Snacktivity, the SnackApp was developed to integrate a range of BCTs that used different app elements (ie, push notifications, data, graphs, and a planner for activity snacks; Table 1).

Snacktivity and Physical Activity Tracking

For users to self-monitor their goals and receive feedback about their Snacktivity, decisions needed to be made about how physical activity should be measured and how users’ progress from the SnackApp was presented and fed back to facilitate their engagement with Snacktivity. One way to achieve this is by using device-based measures of physical activity (eg, accelerometers, fitness trackers, and smartwatches) that can record physical activity continuously. Among the many different consumer wearable physical activity monitors (based on the findings from step 1), the Fitbit Versa smartwatch (Fitbit Inc) was selected. Fitbit Versa has onboard computing that allows data to be analyzed instantaneously to ensure immediate access, feedback, and rewards to users regarding the completion of their Snacktivity goals to promote and facilitate continued engagement with the SnackApp.

SnackApp: Mobile App Development

The research team, working with an app development company, created a series of designs on how the BCTs would be incorporated into SnackApp. These designs were based on modern design techniques (user experience) and previously reviewed apps from commercial and research spaces. These designs were piloted among the planning groups and the participant advisory group and were discussed until a consensus was reached on the best options available.

The SnackApp (Figure 2) is divided into five main areas as follows:

1. “Dashboard”—the main screen of the SnackApp where users can self-monitor their daily activity metrics and progression toward their daily goal
2. “My Stats”—A historical look at the user’s activity to provide feedback on their metrics across days, weeks, and months, also with progression toward their goals
3. “My Goals”—users can set their activity goals relating to activity snacks, steps, and active minutes. Within this area, users can also set action plans related to their goals. Action plans are structures to guide and encourage users to identify when, what, where, and how long the desired behavior change will be conducted (eg, when I am brushing my teeth, I will perform squats for 2 minutes).
4. “My Resources”—this area contains useful examples of activity snacks and how they can be performed safely, articles on the importance of Snacktivity and being physically active, and frequently asked questions regarding the use of the SnackApp and the activity monitor. A “forum” section was included in this area where users of the SnackApp can send messages to each other to discuss Snacktivity and provide social support to engage in Snacktivity.

5. “Notifications”—these formed 2 types: SnackApp and clockface notifications. These notifications were used to provide prompts, cues, or nudges to encourage users to be active.
 - App push notifications: push notifications were grouped into 3 main types: relapse prevention, motivational, and informational; they were prioritized to be delivered in that order. Relapse prevention notifications were designed to guide people in using the app. Motivational messages were designed to help people reach their goals, congratulate participants when they have reached their goals, or encourage them if they have failed to meet their daily targets. Informational messages would be used to tell users that the battery on their watch was running low to ensure that they were charged and therefore in use.
 - Clockface notification: in addition to the push notifications sent to the phone, nudges or prompts could also be displayed on the clockface. After a period of inactivity, the watch would display a message to encourage users to perform an activity snack. If the user was to perform an activity snack within 2 minutes of this prompt, a congratulatory message would appear ([Multimedia Appendix 1](#) provides further information).

There was an aspect of customization to these prompts, as participants could personalize the duration of time before an inactivity prompt was provided (maximum 90 min), as well as the time frame in the day when the SnackApp would send prompts (eg, between 9 AM and 5 PM). These customizations meant that the users were only being nudged in the way decided by them to guard against frustration and disinterest with the SnackApp.

Figure 2. SnackApp and Fitbit clockface.

Calculation of Activity Snacks

Activity snacks are defined as short bouts of MVPA lasting 2 to 5 minutes. Fitbit uses the Karvonen formula to determine activity intensity and returns this as active minutes within their system. Therefore, 2 to 5 active minutes would be considered an activity snack.

Ethics Approval and Informed Consent

This study received favorable ethical opinion from the London City and East Research Ethics Committee (20/PR/0323). Eligible participants were sent a participant information sheet, and they provided written informed consent before any data collection took place.

User Testing of the SnackApp

The next stage of the IM process was to obtain user feedback on SnackApp to allow further development and refinement.

Design, Setting, and Participants

Physically inactive adults who had completed a previous work package within the Snackivity program [33] were recruited to test the first version of SnackApp. Participants were defined as inactive based on their responses to the General Practice physical activity questionnaire. Those who did not use a commercial physical activity tracker but did own an Apple or Android smartphone compatible with iOS 10+ or Android 4.1+ were recruited.

Data Collection

Once participants had provided informed consent, they were sent the Fitbit Versa Lite via post. Participants were also sent instructions on how to download, install, sign in, and connect

to the SnackApp and Fitbit app and were asked to incorporate Snackivity into their daily lives for 28 consecutive days while wearing the Fitbit (with the Snackivity watch face) and using the SnackApp. All data were anonymized before analysis.

Measures

User Version of the Mobile Application Rating Scale

Participants were asked to complete the user version of the mobile application rating scale (uMARS [39]) and report their answers specifically concerning their experiences of using the SnackApp at the end of the data collection period. There are 3 components assessed by the uMARS: “App Quality” (engagement, functionality, esthetics, and information); “App Subjective Quality” (whether participants would recommend the app to others); and perceived impact (whether the app has increased the participant’s awareness, knowledge, attitudes, intention to change, help seeking, and behavior change).

User Engagement

Engagement with consumer physical activity trackers and mHealth apps is not well defined in the literature [40], and there is no standardized list of metrics that can be described and measured for engagement with mHealth apps. Therefore, we defined user engagement as the participants’ use and interactions, captured directly via the SnackApp. Furthermore, the “Display on/off” time (ie, the amount of time the Fitbits screen was on or off henceforth called “Clockface queries”) was measured to provide a proxy indication of the time spent interacting at the SnackApp clockface.

Data Analyses

User engagement was collated and analyzed to provide data on the number of bouts (per day) and the duration of time that users accessed the SnackApp, as well as the time spent on each of the specific screens within the SnackApp. Furthermore, the number and duration of time users viewed the SnackApp clockface as displayed on the Fitbit were summarized, and any bout duration of “clockface display on” >5 seconds was used as a proxy for engagement with the SnackApp information displayed on the Fitbit; 5 seconds was selected for pragmatic reasons because it was considered the minimum time when people could be reasonably assumed to be looking at the SnackApp clockface and to distinguish from the time people might be only determining the time of day, which typically takes only 1 to 2 seconds.

A Sankey analysis diagram was generated using the SnackApp user engagement data. Sankey diagrams (the outcome of a Sankey analysis) are visualizations used to depict the flow from one set of values to another. Sankey analysis are best used in mobile analytics to show “traffic” flows from one app page to another. Each link between the “node” represents a change in the screen viewed in the SnackApp, whereas each “node” represents a particular screen within the SnackApp.

Data relating to user engagement with SnackApp and SnackApp tracker watch clockface on the Fitbit were downloaded from bespoke customer relationship management software and analyzed using proprietary software (The SnackApp Usage package) developed in R Statistical Computing Language (R Foundation for Statistical Computing), which was further developed into a Shiny application [41,42]. The source code for the app (SnackApp Usage package version 0.9.2) can be found on GitHub [43,44].

The subscale and total scores for the uMARS were calculated. The total quality score of the SnackApp was determined by averaging the mean scores of the engagement, functionality, esthetics, and information subscales. The app’s subjective quality subscale items were reported individually. Given the purpose of this research, it was not appropriate to conduct statistical means difference or relationship testing; instead, summary data are presented.

Results

Participants

A total of 15 adults (female, n=8, 53%; male, n=7, 47%; mean age: mean 53, SD 15.7 years; mean self-reported BMI 27.3, SD 4.2 kg/m²) took part in the study (Table 2).

Table 2. Participant demographics (N=15).

	Values
Age (years), mean (SD)	53.0 (15.7)
BMI (kg/m ²), mean (SD)	27.3 (4.2)
Sex, n (%)	
Male	7 (47)
Female	8 (53)
Self-reported ethnicity, n (%)	
White	10 (67)
Black	2 (13)
Pakistani	3 (20)

Engagement

SnackApp

Over the 28-day study period, participants engaged with the SnackApp on average 77 (SD 80) times, for an average total SnackApp time use of 12.6 (SD 47) minutes per week, with most of this time spent on the SnackApp dashboard (14, SD 12.1 bouts lasting 7.5, SD 31 min/week). On average, engagement with the SnackApp was higher among male individuals than among female when considering the overall use (male=24 instances, 15 min/week vs female=15 instances, 11 min/week) and all individual app elements, except for “my goals” (male=2 instances, 1.5 min/week; female=3 instances, 0.6 min/week) and “activity planner” (male=0.2 instances, 0.6 min; female=2 instances, 0.6 min/week; Figure 3).

Figure 4 shows a Sankey diagram, which illustrates that most SnackApp interactions began on the dashboard (default landing page) and typically progressed to the “my stats” and “my goals” pages. Users, who did not initially interact via the dashboard, typically engaged with the dashboard as the second page. The “activity planner” and “forum” pages tended to be the third or fourth pages engaged with on the SnackApp. The height of the column on the Sankey diagram (Figure 3) indicates the number of users who interacted with the app in that order. For example, column 5 is shorter than column one, as fewer users made ≥5 interactions with the SnackApp. The Sankey diagram shows that there are a diminishing number of users interacting with the app after the fifth column, suggesting that most users performed 5 interactions with the app before exiting the app.

Figure 3. Average time spent (minutes) and number of instances on the SnackApp and individual elements of the SnackApp per week.





Figure 4. Sankey diagram representing the flow of participants through the SnackApp from the first page on opening the app through to the 10th page subsequently. Red: dashboard; gray: FAQ; yellow: my goals; green: my stats; purple: activity planner; blue: profile; and amber: forum. FAQ: frequently asked questions. Code available on Github.



Physical Activity Monitor (Fitbit)

Participant SnackApp clockface was queried 112 times per day, of which 52 queries were sustained for >5 seconds

(accumulating a median time of 8.4 min spent querying the clockface per day). Almost 80% of the total time querying the SnackApp clockface display was accounted for in the sustained queries (Table 3).

Table 3. Fitbit and SnackApp clockface queries.

	Values, median (range)
Daily clockface queries (frequency per day)	111.7 (34-285)
Male	97.4 (34-152)
Female	153.9 (60-285)
Sustained^a clockface queries (number per day)	51.8 (23-105)
Male	40.0 (34-152)
Female	64.7 (60-285)
Duration (sustained minutes^a or total min/day)	8.4 (3.6-28.9)
Male	8.2 (3.8-9.2)
Female	11.1 (3.6-28.9)

^aSustained queries are defined as the clockface being on for 5 seconds or longer.

uMARS Score

The mean overall app quality score was 3.5 (SD 0.6) out of 5, with the highest mean component score for esthetics and information (mean score 3.6). Results from the perceived impact

subscale showed that participants “agreed” that the SnackApp increased their awareness, knowledge, and attitude toward increasing physical activity as well as their intention to change and behavior change (Table 4).

Table 4. User version of the mobile application rating scale results from the SnackApp.

	Mean score (SD; out of 5)	Score definition
App quality score	3.5 (0.6)	— ^a
Engagement	3.4 (0.7)	—
Functionality	3.3 (0.7)	—
Esthetics	3.6 (0.6)	—
Information	3.6 (1.0)	—
App subjective quality		
Would you recommend this app to people who might benefit from it?	3 (1.4)	Maybe
How many times would you use the app in the next 12 months use?	4 (1.2)	10-50
Would you pay for this app?	1 (0.5)	Maybe
What is your overall star rating of the app?	3.5 (0.5)	Average
Perceived impact		
Awareness	4.2 (1.4)	Agree
Knowledge	3.8 (1.4)	Agree
Attitudes	4.1 (1.4)	Agree
Intention to change	4.0 (1.2)	Agree
Behavior change	4.2 (0.9)	Agree

^aNot available.

Discussion

IM Framework

This report presented a systematic method that combines conceptual and technological frameworks in the development of a new health app called SnackApp, which will contribute to the future development of the Snacktivity intervention. Specifically, we have described the practical application of the IM framework to the development of Snacktivity intervention

by identifying the behaviors of interest (ie, main behavioral facilitators of physical activity and sedentary behavior) and the application of specific BCTs to bring about health behavior change using mHealth technologies.

User Testing

User testing of the SnackApp focused on utility and engagement, intending to increase users’ enjoyment and motivation of the app [45,46]. Our approach provided an understanding of how participants used and moved through the SnackApp, such that

the intended outcome of the behavior change intervention (eg, completing activity snacks) was optimized.

The number of times users need to engage with mHealth interventions for effective behavior change to occur, across different populations, contexts, and time frames, is unknown. Our data indicated that users engaged with the individual elements of the SnackApp to varying degrees, with the dashboard, stats pages, and goal-setting elements being those that participants interacted with the most. This is encouraging, given that these components of the app encompass the BCTs of goal setting, feedback on progression toward goals, and self-monitoring, which previous research has shown as key ingredients in facilitating physical activity behavior changes [20]. The average total SnackApp time was 12.6 minutes per week. It is currently unknown what level of engagement with digital health intervention would be needed to initiate and sustain behavior change, but participants would likely need to engage with the SnackApp more, in the short term, to sustain any behavior change and to form habits. That said, Yardley et al [47] argued that paradoxically, as habits are formed, this may result in lower use of the SnackApp and users become less reliant on the intervention for behavior change.

Ratings of subjective app quality indicated that participants considered the SnackApp to be of fair to good quality. Although the average engagement score of uMARS was 3.5 out of 5, users indicated that they did not find SnackApp entertaining. The “entertainment” aspect of the “app engagement” score was 3 out of 5 (which suggests that the app was “OK, fun enough to entertain user for a brief time [<5 minutes]”). While this is only a modest rating of the app’s entertainment, entertainment is not the main function of SnackApp. Implementing or increasing enjoyment when using health apps is difficult because enjoyment is not their primary role. However, other health apps have included gamification opportunities, often through avatars, unlocking of areas or levels within the app after a certain period or as a reward for progression and competition among users. It may be that efforts should be made as part of future iterations of SnackApp to include such features.

Strengths and Limitations

This study describes in detail the key perspectives of IM that informed the development of SnackApp through the combination of evidence; participation and engagement from patients, public, and academic stakeholders; and the use of behavioral change theories and BCTs. Overall, our development work was based on a well-established and comprehensive theoretical framework. It is also important to highlight that although the needs assessment and evidence synthesis used here were based predominantly on findings from systematic reviews conducted previously, our underpinning evidence of mHealth use for health behavior change was also based on rigorous studies that had evaluated digital programs.

Several technological strengths relating to the SnackApp merit further discussion. The SnackApp has been designed using the React Native platform (Meta Platforms Inc). This platform enables the production of 1 app across multiple mobile platforms (eg, iOS and Android), meaning the app can reach a larger population. In addition, it is cheaper to develop (and

subsequently maintain and update) 1 code base rather than 2 separate code bases for each platform. Furthermore, the use of the Fitbit Versa smartwatch series provides the potential for data to be collected, analyzed, and fed back to participants, increasing the capability for immediate feedback on Snackactivity behavior to users. This is important because Snackactivity is an approach that is likely to be most potent when participants engage with it throughout the whole day, so the need for instantaneous feedback is beneficial. In addition, the SnackApp system collected comprehensive data about the engagement of users within the SnackApp and with the Fitbit clockface. In the future, these data will be important for understanding how users engage with Snackactivity to understand any mediators to participation in Snackactivity™.

This study has some limitations. The study analysis focused on how and whether users would engage with the SnackApp, but as participants were only followed for 28 days, it is difficult to determine the level of attrition over a longer period (eg, 1 year). Indeed, within the mobile app industry, many engagement and attrition metrics are calculated over several months of use rather than 1 month. Future studies should assess the feasibility and effectiveness of such interventions. Furthermore, SnackApp currently connects only to Fitbit Versa series devices. This will affect the ability of SnackApp to scale up the population level. Future iterations of SnackApp should integrate it across a wider range of wearable devices.

Future Research and Implications

The IM framework is a useful resource that helped to develop a robust digital self-monitoring program for our target audience. Future studies should consider assessing the feasibility and effectiveness of SnackApp and accompanying SnackApp watch face over a longer period.

Although the behavioral goal setting was individually and dynamically tailored to each participant’s physical activity behavior patterns, tailoring may be enhanced by considering additional variables, such as age, sex, health literacy, self-efficacy, intention, and environmental factors (eg, weather patterns), which are likely to affect physical activity behavior and may lead to higher personal relevance of the SnackApp to participants and subsequently contribute to long-term behavior change [48–50]. Furthermore, our user testing was on a small population with a vested interest in Snackactivity; therefore, it would be useful to assess the usability of SnackApp in a larger, more diverse population. As part of future research, a qualitative study could seek to understand the experiences (thoughts and feelings) participants had while using the SnackApp, which would support its further development.

The results of user testing and feedback will be used to inform modifications to the SnackApp. For example, feedback suggests that there is a need to increase the enjoyment of the app to help maintain engagement. This can be achieved through the gamification of the app. Furthermore, clear messaging could be provided in the app on how to complete an activity snack in terms of the intensity and duration of activities. Furthermore, SnackApp currently uses Fitbit to measure participants’

Snacktivity. Future research should look to specifically validate Fitbit as a measure of short bouts of physical activity.

Conclusions

The IM framework was used to aid the development of a theory- and evidence-based mHealth intervention called Snacktivity. IM contributed to the identification of the determinants and

optimal theoretical methods to promote participation in Snacktivity. User testing was also conducted, suggesting that physically inactive adults will engage with the SnackApp. Future research needs to be conducted to assess how engagement with SnackApp will change over time, as well as contribute to intervention effectiveness.

Acknowledgments

Snacktivity study team includes authors AJD, DE, KG, CJG, RM, JS, and LBS, and the contributors Stuart Biddle, Charlotte Edwardson, Emma Frew, Sheila M Greenfield, Natalie Ives, Kate Jolly, Nanette Mutrie, Helen M Parretti, Magdalena Skrybant, and Tom Yates. The authors would like to thank the public advisory group who contributed to this research. The authors would like to thank the Birmingham Community Health care Foundation National Health Service Trust, the Clinical Research Network, West Midlands, and the general practices who supported the study. The authors would like to thank all those who participated in this research. This project was funded as part of the National Institute for Health Research (NIHR) under its Program Grants for Applied Research (RP-PG-0618-20008). AJD was supported by the NIHR Research Professorship award. This research was supported by the NIHR Leicester Biomedical Research Centre. The views expressed are those of the authors and not necessarily those of the National Health Service, NIHR, or Department of Health and Social Care.

Data and Code Availability

The data sets generated and analyzed during this study are available from the corresponding author on reasonable request. Software application or custom code is available on the internet [44].

Authors' Contributions

AJD conceived the concept of Snacktivity. JS collected data and conceived the study. JS and JJCT conducted the analysis. JJCT developed the Shiny app used for analysis. JS, KG, JCR, LBS, DE, RM, AJD, and the Snacktivity study team were involved in the planning and development of SnackApp and provided input and critical feedback on the manuscript. The Snacktivity study team provided feedback at a later stage.

Conflicts of Interest

None declared.

Multimedia Appendix 1

SnackApp technical developments: additional information.

[\[DOCX File , 25 KB-Multimedia Appendix 1\]](#)

References

1. Lee I, Shiroma EJ, Lobelo F, Puska P, Blair SN, Katzmarzyk PT, Lancet Physical Activity Series Working Group. Effect of physical inactivity on major non-communicable diseases worldwide: an analysis of burden of disease and life expectancy. *Lancet* 2012 Jul 21;380(9838):219-229 [FREE Full text] [doi: [10.1016/S0140-6736\(12\)61031-9](https://doi.org/10.1016/S0140-6736(12)61031-9)] [Medline: [22818936](https://pubmed.ncbi.nlm.nih.gov/22818936/)]
2. Kohl HW, Craig CL, Lambert EV, Inoue S, Alkandari JR, Leetongin G, Lancet Physical Activity Series Working Group. The pandemic of physical inactivity: global action for public health. *Lancet* 2012 Jul 21;380(9838):294-305 [FREE Full text] [doi: [10.1016/S0140-6736\(12\)60898-8](https://doi.org/10.1016/S0140-6736(12)60898-8)] [Medline: [22818941](https://pubmed.ncbi.nlm.nih.gov/22818941/)]
3. Wilmot EG, Edwardson CL, Achana FA, Davies MJ, Gorely T, Gray LJ, et al. Sedentary time in adults and the association with diabetes, cardiovascular disease and death: systematic review and meta-analysis. *Diabetologia* 2012 Nov;55(11):2895-2905. [doi: [10.1007/s00125-012-2677-z](https://doi.org/10.1007/s00125-012-2677-z)] [Medline: [22890825](https://pubmed.ncbi.nlm.nih.gov/22890825/)]
4. Greaves CJ, Sheppard KE, Abraham C, Hardeman W, Roden M, Evans PH, IMAGE Study Group. Systematic review of reviews of intervention components associated with increased effectiveness in dietary and physical activity interventions. *BMC Public Health* 2011 Feb 18;11:119 [FREE Full text] [doi: [10.1186/1471-2458-11-119](https://doi.org/10.1186/1471-2458-11-119)] [Medline: [21333011](https://pubmed.ncbi.nlm.nih.gov/21333011/)]
5. Samdal GB, Eide GE, Barth T, Williams G, Meland E. Effective behaviour change techniques for physical activity and healthy eating in overweight and obese adults; systematic review and meta-regression analyses. *Int J Behav Nutr Phys Act* 2017 Mar 28;14(1):42 [FREE Full text] [doi: [10.1186/s12966-017-0494-y](https://doi.org/10.1186/s12966-017-0494-y)] [Medline: [28351367](https://pubmed.ncbi.nlm.nih.gov/28351367/)]
6. UK chief medical officers' physical activity guidelines. Department of Health & Social Care. 2019. URL: https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/832868/uk-chief-medical-officers-physical-activity-guidelines.pdf [accessed 2022-08-18]

7. Bull FC, Al-Ansari SS, Biddle S, Borodulin K, Buman MP, Cardon G, et al. World Health Organization 2020 guidelines on physical activity and sedentary behaviour. *Br J Sports Med* 2020 Dec;54(24):1451-1462 [[FREE Full text](#)] [doi: [10.1136/bjsports-2020-102955](https://doi.org/10.1136/bjsports-2020-102955)] [Medline: [33239350](https://pubmed.ncbi.nlm.nih.gov/33239350/)]
8. Hillsdon M, Foster C, Thorogood M. Interventions for promoting physical activity. *Cochrane Database Syst Rev* 2005 Jan 25(1):CD003180 [[FREE Full text](#)] [doi: [10.1002/14651858.CD003180.pub2](https://doi.org/10.1002/14651858.CD003180.pub2)] [Medline: [15674903](https://pubmed.ncbi.nlm.nih.gov/15674903/)]
9. Kwasnicka D, Dombrowski SU, White M, Sniehotta F. Theoretical explanations for maintenance of behaviour change: a systematic review of behaviour theories. *Health Psychol Rev* 2016 Sep;10(3):277-296 [[FREE Full text](#)] [doi: [10.1080/17437199.2016.1151372](https://doi.org/10.1080/17437199.2016.1151372)] [Medline: [26854092](https://pubmed.ncbi.nlm.nih.gov/26854092/)]
10. Zenko Z, Willis EA, White DA. Proportion of adults meeting the 2018 physical activity guidelines for Americans according to accelerometers. *Front Public Health* 2019 Jun 7;7:135 [[FREE Full text](#)] [doi: [10.3389/fpubh.2019.00135](https://doi.org/10.3389/fpubh.2019.00135)] [Medline: [31231627](https://pubmed.ncbi.nlm.nih.gov/31231627/)]
11. Bennie JA, Kolbe-Alexander T, Seghers J, Biddle SJ, De Cocker K. Trends in muscle-strengthening exercise among nationally representative samples of United States adults between 2011 and 2017. *J Phys Act Health* 2020 May 01;17(5):512-518. [doi: [10.1123/jpah.2019-0472](https://doi.org/10.1123/jpah.2019-0472)] [Medline: [32283540](https://pubmed.ncbi.nlm.nih.gov/32283540/)]
12. Bennie JA, De Cocker K, Smith JJ, Wiesner GH. The epidemiology of muscle-strengthening exercise in Europe: a 28-country comparison including 280,605 adults. *PLoS One* 2020;15(11):e0242220 [[FREE Full text](#)] [doi: [10.1371/journal.pone.0242220](https://doi.org/10.1371/journal.pone.0242220)] [Medline: [33237930](https://pubmed.ncbi.nlm.nih.gov/33237930/)]
13. Physical Activity Guidelines Advisory Committee Scientific Report. U.S. Department of Health and Human Services. 2018. URL: https://health.gov/sites/default/files/2019-09/PAG_Advisory_Committee_Report.pdf [accessed 2022-08-18]
14. Biswas A, Oh PI, Faulkner GE, Bajaj RR, Silver MA, Mitchell MS, et al. Sedentary time and its association with risk for disease incidence, mortality, and hospitalization in adults: a systematic review and meta-analysis. *Ann Intern Med* 2015 Jan 20;162(2):123-132. [doi: [10.7326/M14-1651](https://doi.org/10.7326/M14-1651)] [Medline: [25599350](https://pubmed.ncbi.nlm.nih.gov/25599350/)]
15. Ekelund U, Tarp J, Steene-Johannessen J, Hansen BH, Jefferis B, Fagerland MW, et al. Dose-response associations between accelerometry measured physical activity and sedentary time and all cause mortality: systematic review and harmonised meta-analysis. *BMJ* 2019 Aug 21;366:l4570 [[FREE Full text](#)] [doi: [10.1136/bmj.l4570](https://doi.org/10.1136/bmj.l4570)] [Medline: [31434697](https://pubmed.ncbi.nlm.nih.gov/31434697/)]
16. Sanders J, Biddle S, Gokal K, Sherar L, Skrybant M, Parretti H, Snackitivity Study Team. 'Snackitivity™' to increase physical activity: time to try something different? *Prev Med* 2021 Dec;153:106851. [doi: [10.1016/j.ypmed.2021.106851](https://doi.org/10.1016/j.ypmed.2021.106851)] [Medline: [34662595](https://pubmed.ncbi.nlm.nih.gov/34662595/)]
17. Stamatakis E, Huang B, Maher C, Thøgersen-Ntoumani C, Stathi A, Dempsey PC, et al. Untapping the health enhancing potential of vigorous intermittent lifestyle physical activity (VILPA): rationale, scoping review, and a 4-pillar research framework. *Sports Med* 2021 Jan;51(1):1-10 [[FREE Full text](#)] [doi: [10.1007/s40279-020-01368-8](https://doi.org/10.1007/s40279-020-01368-8)] [Medline: [33108651](https://pubmed.ncbi.nlm.nih.gov/33108651/)]
18. Marcolino MS, Oliveira JA, D'Agostino M, Ribeiro AL, Alkmim MB, Novillo-Ortiz D. The impact of mHealth interventions: systematic review of systematic reviews. *JMIR Mhealth Uhealth* 2018 Jan 17;6(1):e23 [[FREE Full text](#)] [doi: [10.2196/mhealth.8873](https://doi.org/10.2196/mhealth.8873)] [Medline: [29343463](https://pubmed.ncbi.nlm.nih.gov/29343463/)]
19. Romeo A, Edney S, Plotnikoff R, Curtis R, Ryan J, Sanders I, et al. Can smartphone apps increase physical activity? Systematic review and meta-analysis. *J Med Internet Res* 2019 Mar 19;21(3):e12053 [[FREE Full text](#)] [doi: [10.2196/12053](https://doi.org/10.2196/12053)] [Medline: [30888321](https://pubmed.ncbi.nlm.nih.gov/30888321/)]
20. Michie S, Abraham C, Whittington C, McAteer J, Gupta S. Effective techniques in healthy eating and physical activity interventions: a meta-regression. *Health Psychol* 2009 Nov;28(6):690-701. [doi: [10.1037/a0016136](https://doi.org/10.1037/a0016136)] [Medline: [19916637](https://pubmed.ncbi.nlm.nih.gov/19916637/)]
21. Aromatario O, Van Hove A, Vuillemin A, Foucaut A, Crozet C, Pommier J, et al. How do mobile health applications support behaviour changes? A scoping review of mobile health applications relating to physical activity and eating behaviours. *Public Health* 2019 Oct;175:8-18 [[FREE Full text](#)] [doi: [10.1016/j.puhe.2019.06.011](https://doi.org/10.1016/j.puhe.2019.06.011)] [Medline: [31374453](https://pubmed.ncbi.nlm.nih.gov/31374453/)]
22. Afshin A, Babalola D, Mclean M, Yu Z, Ma W, Chen C, et al. Information technology and lifestyle: a systematic evaluation of internet and mobile interventions for improving diet, physical activity, obesity, tobacco, and alcohol use. *J Am Heart Assoc* 2016 Aug 31;5(9):e003058 [[FREE Full text](#)] [doi: [10.1161/JAHA.115.003058](https://doi.org/10.1161/JAHA.115.003058)] [Medline: [27581172](https://pubmed.ncbi.nlm.nih.gov/27581172/)]
23. Lunde P, Nilsson BB, Bergland A, Kværner KJ, Bye A. The effectiveness of smartphone apps for lifestyle improvement in noncommunicable diseases: systematic review and meta-analyses. *J Med Internet Res* 2018 May 04;20(5):e162 [[FREE Full text](#)] [doi: [10.2196/jmir.9751](https://doi.org/10.2196/jmir.9751)] [Medline: [29728346](https://pubmed.ncbi.nlm.nih.gov/29728346/)]
24. Muellmann S, Forberger S, Möllers T, Bröring E, Zeeb H, Pischke CR. Effectiveness of eHealth interventions for the promotion of physical activity in older adults: a systematic review. *Prev Med* 2018 Mar;108:93-110. [doi: [10.1016/j.ypmed.2017.12.026](https://doi.org/10.1016/j.ypmed.2017.12.026)] [Medline: [29289643](https://pubmed.ncbi.nlm.nih.gov/29289643/)]
25. Craig P, Dieppe P, Macintyre S, Michie S, Nazareth I, Petticrew M, Medical Research Council Guidance. Developing and evaluating complex interventions: the new Medical Research Council guidance. *BMJ* 2008 Sep 29;337:a1655 [[FREE Full text](#)] [doi: [10.1136/bmj.a1655](https://doi.org/10.1136/bmj.a1655)] [Medline: [18824488](https://pubmed.ncbi.nlm.nih.gov/18824488/)]
26. Bartholome E, Parcel G, Kok G, Gottlieb N, Fernandez M, Eldredge L. *Planning Health Promotion Programs An Intervention Mapping Approach*. Hoboken, New Jersey: Wiley; 2011.
27. Skivington K, Matthews L, Simpson SA, Craig P, Baird J, Blazeby JM, et al. A new framework for developing and evaluating complex interventions: update of Medical Research Council guidance. *BMJ* 2021 Sep 30;374:n2061 [[FREE Full text](#)] [doi: [10.1136/bmj.n2061](https://doi.org/10.1136/bmj.n2061)] [Medline: [34593508](https://pubmed.ncbi.nlm.nih.gov/34593508/)]

28. Daley AJ, Griffin RA, Moakes CA, Sanders JP, Skrybant M, Ives N, et al. Snackivity™ to promote physical activity and reduce future risk of disease in the population: protocol for a feasibility randomised controlled trial and nested qualitative study. *Pilot Feasibility Stud* 2023 Mar 17;9(1):45-95 [FREE Full text] [doi: [10.1186/s40814-023-01272-8](https://doi.org/10.1186/s40814-023-01272-8)] [Medline: [36932423](https://pubmed.ncbi.nlm.nih.gov/36932423/)]
29. Peddie MC, Bone JL, Rehrer NJ, Skeaff CM, Gray AR, Perry TL. Breaking prolonged sitting reduces postprandial glycemia in healthy, normal-weight adults: a randomized crossover trial. *Am J Clin Nutr* 2013 Aug;98(2):358-366 [FREE Full text] [doi: [10.3945/ajcn.112.051763](https://doi.org/10.3945/ajcn.112.051763)] [Medline: [23803893](https://pubmed.ncbi.nlm.nih.gov/23803893/)]
30. Dunstan DW, Kingwell BA, Larsen R, Healy GN, Cerin E, Hamilton MT, et al. Breaking up prolonged sitting reduces postprandial glucose and insulin responses. *Diabetes Care* 2012 May;35(5):976-983 [FREE Full text] [doi: [10.2337/dc11-1931](https://doi.org/10.2337/dc11-1931)] [Medline: [22374636](https://pubmed.ncbi.nlm.nih.gov/22374636/)]
31. Benatti FB, Ried-Larsen M. The effects of breaking up prolonged sitting time: a review of experimental studies. *Med Sci Sports Exerc* 2015 Oct;47(10):2053-2061. [doi: [10.1249/MSS.0000000000000654](https://doi.org/10.1249/MSS.0000000000000654)] [Medline: [26378942](https://pubmed.ncbi.nlm.nih.gov/26378942/)]
32. Going the extra mile: improving the nation's health and wellbeing through public involvement in research. National Institute for Health Research. URL: <https://www.nihr.ac.uk/documents/about-us/our-contribution-to-research/how-we-involve-patients-carers-and-the-public/Going-the-Extra-Mile.pdf> [accessed 2022-08-18]
33. Gokal K, Amos-Hirst R, Moakes CA, Sanders JP, Eslinger DW, Sherar LB, et al. Views of the public about Snackivity™: a small changes approach to promoting physical activity and reducing sedentary behaviour. *BMC Public Health* 2022 Mar 29;22(1):618 [FREE Full text] [doi: [10.1186/s12889-022-13050-x](https://doi.org/10.1186/s12889-022-13050-x)] [Medline: [35351075](https://pubmed.ncbi.nlm.nih.gov/35351075/)]
34. Michie S, Richardson M, Johnston M, Abraham C, Francis J, Hardeman W, et al. The behavior change technique taxonomy (v1) of 93 hierarchically clustered techniques: building an international consensus for the reporting of behavior change interventions. *Ann Behav Med* 2013 Aug;46(1):81-95 [FREE Full text] [doi: [10.1007/s12160-013-9486-6](https://doi.org/10.1007/s12160-013-9486-6)] [Medline: [23512568](https://pubmed.ncbi.nlm.nih.gov/23512568/)]
35. Carver CS, Scheier MF. Control theory: a useful conceptual framework for personality-social, clinical, and health psychology. *Psychol Bull* 1982 Jul;92(1):111-135. [Medline: [7134324](https://pubmed.ncbi.nlm.nih.gov/7134324/)]
36. Kanfer F, Gaelick-Buys L. Self-management methods. In: *Helping People Change: A Textbook of Methods*. Oxford, United Kingdom: Pergamon Press; 1991.
37. Deci E, Ryan R. *Intrinsic Motivation and Self-Determination in Human Behavior*. United States: Springer; 1975.
38. Lally P, Gardner B. Promoting habit formation. *Health Psychol Rev* 2013 May;7(sup1):S137-S158. [doi: [10.1080/17437199.2011.603640](https://doi.org/10.1080/17437199.2011.603640)]
39. Stoyanov SR, Hides L, Kavanagh DJ, Wilson H. Development and validation of the user version of the mobile application rating scale (uMARS). *JMIR Mhealth Uhealth* 2016 Jun 10;4(2):e72 [FREE Full text] [doi: [10.2196/mhealth.5849](https://doi.org/10.2196/mhealth.5849)] [Medline: [27287964](https://pubmed.ncbi.nlm.nih.gov/27287964/)]
40. Arigo D, Jake-Schoffman DE, Wolin K, Beckjord E, Hekler EB, Pagoto SL. The history and future of digital health in the field of behavioral medicine. *J Behav Med* 2019 Feb;42(1):67-83 [FREE Full text] [doi: [10.1007/s10865-018-9966-z](https://doi.org/10.1007/s10865-018-9966-z)] [Medline: [30825090](https://pubmed.ncbi.nlm.nih.gov/30825090/)]
41. Chang W, Cheng J, Allaire J, Sievert C, Schloerke B, Xie Y, SpryMedia Limited, jQuery UI contributors, jQuery Foundation, R Core Team. shiny: Web Application Framework for R. CRAN R Project. 2022. URL: <https://cran.r-project.org/web/packages/shiny/index.html> [accessed 2022-08-18]
42. SnackApp Usage Data Upload. SnackApp. URL: <https://climb.shinyapps.io/snackapp/> [accessed 2022-04-18]
43. R: a language and environment for statistical computing. GBIF. URL: <https://www.gbif.org/tool/81287/r-a-language-and-environment-for-statistical-computing> [accessed 2022-01-31]
44. SnackApp usage. Github. URL: <https://github.com/lboro-climb/snackapp.usage> [accessed 2022-04-18]
45. Kim YH, Kim DJ, Wachter K. A study of mobile user engagement (MoEN): engagement motivations, perceived value, satisfaction, and continued engagement intention. *Decis Support Syst* 2013 Dec;56:361-370. [doi: [10.1016/j.dss.2013.07.002](https://doi.org/10.1016/j.dss.2013.07.002)]
46. O'Brien HL. The influence of hedonic and utilitarian motivations on user engagement: the case of online shopping experiences. *Interact Comput* 2010 Sep;22(5):344-352. [doi: [10.1016/j.intcom.2010.04.001](https://doi.org/10.1016/j.intcom.2010.04.001)]
47. Yardley L, Morrison L, Bradbury K, Muller I. The person-based approach to intervention development: application to digital health-related behavior change interventions. *J Med Internet Res* 2015 Jan 30;17(1):e30 [FREE Full text] [doi: [10.2196/jmir.4055](https://doi.org/10.2196/jmir.4055)] [Medline: [25639757](https://pubmed.ncbi.nlm.nih.gov/25639757/)]
48. Morrison LG, Yardley L, Powell J, Michie S. What design features are used in effective e-health interventions? A review using techniques from critical interpretive synthesis. *Telemed J E Health* 2012 Mar;18(2):137-144 [FREE Full text] [doi: [10.1089/tmj.2011.0062](https://doi.org/10.1089/tmj.2011.0062)] [Medline: [22381060](https://pubmed.ncbi.nlm.nih.gov/22381060/)]
49. Head KJ, Noar SM, Iannarino NT, Grant Harrington N. Efficacy of text messaging-based interventions for health promotion: a meta-analysis. *Soc Sci Med* 2013 Nov;97:41-48. [doi: [10.1016/j.socscimed.2013.08.003](https://doi.org/10.1016/j.socscimed.2013.08.003)] [Medline: [24161087](https://pubmed.ncbi.nlm.nih.gov/24161087/)]
50. Hekler EB, Rivera DE, Martin CA, Phatak SS, Freigoun MT, Korinek E, et al. Tutorial for using control systems engineering to optimize adaptive mobile health interventions. *J Med Internet Res* 2018 Jun 28;20(6):e214 [FREE Full text] [doi: [10.2196/jmir.8622](https://doi.org/10.2196/jmir.8622)] [Medline: [29954725](https://pubmed.ncbi.nlm.nih.gov/29954725/)]

Abbreviations

BCT: behavior change technique

IM: intervention mapping

mHealth: mobile health

MoSCoW: Must Have, Should Have, Could Have, and Won't Have

MVPA: moderate- to vigorous-intensity physical activity

PAG: public advisory group

uMARS: user version of the mobile application rating scale

Edited by A Mavragani; submitted 15.07.22; peer-reviewed by H Sediva, J Alvarez Pitti; comments to author 08.12.22; revised version received 31.01.23; accepted 14.02.23; published 22.05.23

Please cite as:

Sanders JP, Gokal K, Thomas JJC, Rawstorn JC, Sherar LB, Maddison R, Greaves CJ, Esliger D, Daley AJ, Snacktivity Investigators. Development of a Mobile Health Snacktivity App to Promote Physical Activity in Inactive Adults (SnackApp): Intervention Mapping and User Testing Study

JMIR Form Res 2023;7:e41114

URL: <https://formative.jmir.org/2023/1/e41114>

doi: [10.2196/41114](https://doi.org/10.2196/41114)

PMID:

©James P Sanders, Kajal Gokal, Jonah J C Thomas, Jonathan C Rawstorn, Lauren B Sherar, Ralph Maddison, Colin J Greaves, Dale Esliger, Amanda J Daley, Snacktivity Investigators. Originally published in JMIR Formative Research (<https://formative.jmir.org>), 22.05.2023. This is an open-access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work, first published in JMIR Formative Research, is properly cited. The complete bibliographic information, a link to the original publication on <https://formative.jmir.org>, as well as this copyright and license information must be included.